



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

7. The parthenitæ are to be looked upon as physiologically young, and thus able to continue the parthenogenetic cycle for several generations without the hermaphrodite generation.

8. This youth and the attendant simplicity as compared with the miracidium is to be looked upon as secondary results of parasitism.

THE CRYSTALLINE STYLE OF LAMELLIBRANCHS

Nelson (Jour. Morp. Vol. 31, June 1918) presents a review of the work done on the crystalline style in Lamellibranchs, and contributes some interesting results of his own about this singular organ.

The lining of the intestine and of the communicating style sac is ciliated. This ciliary mechanism is regarded as having power to separate food from the foreign particles within the tract. Little discrimination is shown as to material taken into the stomach.

The style arises as a thin core of bubbly mucus upon which co-axial layers of a gelatinous protein, containing enzymes, are deposited. The style rotates in the sac, according to the observer. He confirms the conclusion that it contains strong amylolytic ferments and believes that the style serves as a means of restoring to the stomach undigested food particles which might otherwise be lost, at least in those forms in which the style sac is not separated from the intestine. The store of ferment is thought to be of peculiar value because of the long period during which feeding is impossible in many mollusks.

REVIVIFICATION OF EXSICCATED EARTHWORMS

Schmidt (Jour. Exp. Zool., October 1918) has shown that earthworms are capable of being revived after 39 to 48 hours (depending on temperature) of exsiccation in which they have lost one-third to one-half their length and volume, and show no signs of life that can be detected. The worms were of course very gradually dried. The body must be allowed to retain its elasticity and the skin its softness, if revival is to be expected. Life was normally regained after as much as 61.6 per cent of the weight of body (nearly 73 per cent of the weight of water in the body) had been lost.

The earthworms differ from lower animals like rotifers and nematodes in that they cannot be preserved thru such long periods of time. This is probably due to the fact that they are more complicated, cannot be so completely desiccated, and hence decomposition